

## ADJACENCY PROPERTIES OF GRAPHS AND A CONJECTURE OF ERDŐS

ANTHONY BONATO AND ALEXANDRU COSTEA

ABSTRACT. In 1963, Erdős and Rényi gave a non-explicit, randomized construction of graphs with an adjacency property. For a positive integer  $n$ , a graph is *n-exentially closed* (or *n-e.c.*) if for all disjoint sets of vertices  $A$  and  $B$  with  $|A \cup B| = n$ , there is a vertex  $z$  not in  $A \cup B$  joined to each vertex of  $A$  and no vertex of  $B$ . Until recently, only a few explicit families of *n-e.c.* graphs were known, such as Paley graphs. Erdős posed a conjecture on the asymptotic minimum order of *n-e.c.* graphs which is only now receiving attention. The exact minimum orders of *n-e.c.* graphs are only known for  $n = 1$  and  $n = 2$ .

Using a computer search, a new example of a 3-e.c. graph of order 30 is presented. Previously, no known 3-e.c. graph was known to exist of that order. We give a new randomized construction of vertex-transitive *n-e.c.* graphs, exploiting Cayley graphs.

**1. Introduction.** The probabilistic method was discovered by Paul Erdős in the 1950's and has since found widespread use in many fields of Mathematics and Computer Science, and especially so in graph theory. The method shows objects with prescribed properties occur with positive probability, and therefore such objects exist. Finding explicit examples of the objects (in our case, graphs with an adjacency property) is often quite challenging.

An *adjacency property* is a global property of a graph, where given a fixed subset of vertices  $S$ , there exists vertices outside of  $S$  joined to vertices of  $S$  in a predetermined way. Adjacency properties stem from a seminal paper on random graphs by Erdős and Rényi [12] published in 1963. One particular adjacency property that has received much recent attention is the *n-e.c.* property. For a positive integer  $n$ , a graph is *n-exentially closed* (or *n-e.c.*) if for all disjoint sets of vertices  $A$  and  $B$  with  $|A \cup B| = n$  (one of  $A$  or  $B$  can be empty), there is a vertex

---

2000 *Mathematics Subject Classification.* 05C75, 05C80.

*Key words and phrases.* graphs, adjacency property, *n-e.c.* graph, random graphs, vertex-transitive, Cayley graph.

The authors gratefully acknowledge support from NSERC and MITACS.

$z$  not in  $A \cup B$  joined to each vertex of  $A$  and no vertex of  $B$ . We say that  $z$  is *correctly joined* (or *c.j.*) to  $A$  and  $B$ . See Figure 1. Hence,




FIGURE 1. The  $n$ -e.c. property.

for all  $n$ -subsets  $S$  of vertices, there exist  $2^n$ -many vertices joined to  $S$  in all possible ways. Although the  $n$ -e.c. property is straightforward to define, it is not obvious from the definition that graphs with the property exist.

Erdős and Rényi gave a non-explicit, randomized construction of such a graph in [12]. Roughly speaking, random graphs arise by choosing edges among pairs of distinct vertices independently with a given probability. To be more precise, the *random graph*  $G(m, p)$  consists of the probability space  $(\mathcal{G}_m, \mathcal{F}, \mathbb{P})$ , where  $\mathcal{G}_m$  is the set of all graphs with vertex set  $\{1, 2, \dots, m\}$ ,  $\mathcal{F}$  is the family of all subsets of  $\mathcal{G}_m$ . Each graph is chosen independently with probability a fixed  $p \in (0, 1)$ . There are  $|\mathcal{G}_m| = 2^{\binom{m}{2}}$  graphs of order  $m$ , so the probability function is given by

$$\mathbb{P}(G) = p^{|E(G)|} (1 - p)^{\binom{m}{2} - |E(G)|}$$

This space may be viewed as  $\binom{m}{2}$  independent coin flips, one for each pair of vertices where the probability of drawing an edge is equal to  $p$ .

We say that an event holds *asymptotically almost surely* (a.a.s.) in  $G(m, p)$  if it holds with probability tending to 1 as  $m \rightarrow \infty$ . We will consider asymptotic results on probability spaces such as  $G(m, p)$ , so we recall asymptotic notation. Let  $f$  and  $g$  be functions whose domain is some fixed subset of  $\mathbb{R}$ . We write  $f \in O(g)$  if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

exists and is finite. We abuse notation and write  $f = O(g)$ . We write  $f = \Omega(g)$  if  $g = O(f)$ , and  $f = \Theta(g)$  if  $f = O(g)$  and  $f = \Omega(g)$ . If

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0,$$

then  $f = o(g)$ . So if  $f = o(1)$ , then  $f$  tends to 0. We write  $f \sim g$  if  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 1$ . All logarithms are in base  $e$ , and are written as  $\log x$ .

**Theorem 1.** *Fix an integer  $n \in \mathbb{N}$ . The following then holds.*

1. *A.a.s.  $G(m, \frac{1}{2})$  is  $n$ -e.c.*
2. *Let  $f$  be a positive real-valued function defined by*

$$f(m, n) = \binom{m}{n} 2^n \left(1 - \frac{1}{2^n}\right)^{m-n}.$$

*If  $m$  is an integer chosen so that  $f(m, n) < 1$ , then there is an  $n$ -e.c. graph of order  $m$ .*

**Proof.** Let  $G = G(m, 1/2)$ . For item (1), let  $A, B$  be two sets of vertices of  $G$  such that  $A \cap B = \emptyset$  and  $|A \cup B| = n$ . The probability that no vertex of  $G$  is joined correctly to  $A$  and  $B$  is

$$\left(1 - \frac{1}{2^n}\right)^{m-n}. \quad (1)$$

There are  $\binom{m}{n}$  choices of an  $n$ -set of vertices  $X$ , and  $2^n$  many partitions of  $X$  into sets  $A$  and  $B$ . Hence, by (1) it follows that the probability that  $G$  is not  $n$ -e.c. is at most

$$\begin{aligned} \binom{m}{n} 2^n \left(1 - \frac{1}{2^n}\right)^{m-n} &\leq m^n 2^n \left(1 - \frac{1}{2^n}\right)^{m-n} \\ &= \exp(n \log m + n \log 2 + (m-n) \log(1 - \frac{1}{2^n})) \\ &= o(1), \end{aligned}$$

where the last equality follows since  $\log(1 - \frac{1}{2^n})$  is a negative constant.

For the proof of item (2), if  $m$  has the given property, then with positive probability,  $G(m, \frac{1}{2})$  contains a  $n$ -e.c. graph with positive probability.  $\square$

We note that Theorem 1 easily generalizes to  $G(m, p)$ , where  $p \in (0, 1)$  is fixed. We now consider how large  $m$  must be for  $G(m, \frac{1}{2})$  to be  $n$ -e.c. with positive probability. A plot of  $n$  versus  $m$  (where the corresponding  $m$  value is found by solving the equation  $\lceil f(m, n) \rceil = 1$ ) is shown in Figure 2. Note how quickly the function increases with  $n$ .

We now turn our attention to the minimum order of an  $n$ -e.c. graph. For a positive integer  $n$ , denote the minimum order of an  $n$ -e.c. graph by  $m_{ec}(n)$ . By Theorem 1 (2),  $n$ -e.c. graphs exist for all  $n > 0$ , and so the function  $m_{ec}(n)$  is well-defined. It was determined in [8] that  $m_{ec}(1) = 4$  and  $m_{ec}(2) = 9$ . These are the only two known values of this function! In [8], it was shown that there are exactly three non-isomorphic 1-e.c. graphs of order four; see Figure 3.




FIGURE 2. Number of vertices  $m$  needed for  $G(m, 1/2)$  to be  $n$ -e.c.




FIGURE 3. The 1-e.c. graphs of minimum order.

The unique isomorphism type of minimum order 2-e.c. graphs (as proved in [8]) is shown in Figure 4.




FIGURE 4. The unique 2-e.c. graph of minimum order.

We note that Theorem 1 (2) supplies an asymptotic upper bound for  $G(m, \frac{1}{2})$  to be  $n$ -e.c., which we describe in our next theorem. As the proof is part of folklore, but not evident in the literature, so we give it explicitly here.

**Theorem 2.** *If  $m = O(n^2 2^n)$  and  $n$  is a sufficiently large integer, then with positive probability  $G(m, \frac{1}{2})$  is  $n$ -e.c. In particular,*

$$m_{ec}(n) = O(n^2 2^n).$$

**Proof.** We must show that if  $m = O(n^2 2^n)$ , then  $f(m, n) < 1$ . Equivalently, we show that if  $\epsilon > 0$  is fixed and  $m = \lfloor (\epsilon + 1)n^2 2^n \rfloor$ , then

$$\log f(m) < 0. \quad (2)$$

For simplicity, we drop the floor and work with  $m = (\epsilon + 1)n^2 2^n$ . Now

$$\binom{m}{n} 2^n \left(1 - \frac{1}{2^n}\right)^{m-n} < m^n 2^n \left(1 - \frac{1}{2^n}\right)^{m-n}.$$

Hence, (2) is equivalent to showing that

$$n \log m + n \log 2 + (m - n) \log \left(1 - \frac{1}{2^n}\right) < 0. \quad (3)$$

For  $n$  sufficiently large we have that  $\log \left(1 - \frac{1}{2^n}\right) \sim -\frac{1}{2^n}$ . By this fact, by computation, and by the choice of  $m$ , (3) is equivalent to

$$n(\log(\epsilon + 1) + 2 \log n + \log 2) + n^2 \log 2 + \frac{n}{2^n} < (\epsilon + 1)n^2,$$

which is valid for large  $n$  as  $\log 2 < 1$ .  $\square$

The determination of  $m_{ec}(n)$ , where  $n \geq 3$  appears to be an extremely difficult open problem. An asymptotic lower bound was proved in Erdős et al. [10], where it was shown that

$$m_{ec}(n) = \Omega(n 2^n).$$

One of the deepest (and one we think deserves to be better known) conjectures on  $n$ -e.c. graphs was posed by Caccetta, Erdős, and Vijayan [10].

**Conjecture 1.**

$$m_{ec}(n) = \Theta(n 2^n).$$

Hence, to prove the conjecture, we would need to present a family of  $n$ -e.c. graphs with order  $O(n 2^n)$ . Theorem 2 actually gives the best known upper bound to  $m_{ec}(n)$ . We therefore need to reduce the order  $O(n^2 2^n)$  by a multiplicative factor of  $n$ .

There has been much research done in determining the minimum order of a 3-e.c. graph. The results of [8] show that

$$20 \leq m_{ec}(3) \leq 28.$$

A new lower bound on the order of  $m_{ec}(3)$  was found recently using more sophisticated computational methods. Based on 15,000 hours of

CPU time, the authors of [14] demonstrated that  $m_{ec}(3) \geq 24$ . Hence,  $m_{ec}(3)$  can only be one of

$$24, 25, 26, 27, 28.$$

We do not solve Erdős' conjecture here, nor determine  $m_{ec}(3)$ . However, we present some new computational results on 3-e.c. graphs of order 30 in Section 2, and a new construction of vertex-transitive  $n$ -e.c. graphs arising from random Cayley graphs in Section 3. Both results were first presented in the second author's Master's thesis [11].

**1.1. Explicit Constructions of  $n$ -e.c. Graphs.** The first family of explicit graphs that were discovered to contain  $n$ -e.c. graphs for all  $n$  were Paley graphs. Fix  $q$  a prime power with  $q \equiv 1 \pmod{4}$ . A *Paley graph*, written  $P_q$ , is a graph constructed on the points of a finite field of order  $q$  such that two vertices are adjacent if and only if their difference is a non-zero square in the field. As an example, consider the graph  $P_9$ , which is isomorphic to the minimum order 2-e.c. graph. Let  $S$  be the set of all non-zero squares in the finite field with 9 elements, written  $GF(9)$ . We use the following representation of elements of this field:

$$GF(9) = \{a + bi : a, b \in \mathbb{Z}_3, i^2 = -1\}.$$

The non-zero squares are therefore  $\{1, 2, i, 2i\}$ . See Figure 5, where vertices are labelled by the elements of  $GF(9)$ .




FIGURE 5. The graph  $P_9$ .

The following result on the  $n$ -e.c. properties of Paley graphs was proven independently in [4, 6]. The proof uses a famous result from number theory: Weil's proof of the Riemann hypothesis over finite fields.

**Theorem 3.** *If*

$$q > n^2 2^{2n-2},$$

*then  $P_q$  is  $n$ -e.c.*

For a more detailed discussion of construction of  $n$ -e.c. graphs arising from number theory, designs, geometry, and set theory, the reader is directed to the recent survey [7].

**2. Computational results.** The difficulty of determining if a graph is  $n$ -e.c. increases dramatically with  $n$ . For example, to check that a graph of order  $m$  is  $n$ -e.c., for each of the  $\binom{m}{n}$  subsets  $S$  of vertices, we would need to find  $2^n$  vertices joined to  $S$  in all the possible ways. This becomes difficult, if not impossible, to do by hand for large  $n$ . The focus of this section is on computational results related to orders of small 3-e.c. graphs. We recall from the introduction that

$$24 \leq m_{ec}(3) \leq 28$$

(the lower bound follows from [14], while the upper bound follows from [8]). We note that most of the known explicit  $n$ -e.c. graphs are strongly regular. For  $n = 3$ , in [2] it was shown that the Paley graph of order 29 is the minimum order 3-e.c. Paley graph. Few examples of strongly regular non-Paley  $n$ -e.c. graphs are known.

A graph  $G$  is *vertex-transitive* if any two distinct vertices of  $G$ , there is an automorphism mapping one to the other. Hence, in a vertex-transitive graph, any two vertices behave identically. For example, the graph  $P_9$  is vertex-transitive, as is any cycle or clique.

Most of the known explicit  $n$ -e.c. graphs are strongly regular. Let  $k$ ,  $v > 0$ ,  $\lambda$ , and  $\mu$  be non-negative integers. A  $k$ -regular graph  $G$  with  $v$  vertices, so that each pair of joined vertices has exactly  $\lambda$  common neighbours, and each pair of non-joined vertices has exactly  $\mu$  common neighbours, is called a *strongly regular graph*; we say that  $G$  is  $\text{SRG}(v, k, \lambda, \mu)$ . Paley graphs are an important instance of strongly regular graphs; the graph  $P_q$  is a  $\text{SRG}(q, \frac{q-1}{2}, \frac{q-5}{4}, \frac{q-1}{4})$  (for a proof, see [5]). In our computer search for a minimum order 3-e.c. graph, we focused on strongly regular graphs and the class of vertex-transitive graphs with orders between 24 to 30 (inclusive).

Lists containing all isotypes of small order vertex-transitive and strongly regular graphs are publicly available on-line. The data sets for the class of strongly regular graphs can be found in [17], while the data for the class of vertex-transitive graphs is available on-line at [16]. The data set is partitioned into different files based on the order of the graph. Each file consists of adjacency matrices encoded in the *g6* format. (More on this format can be found at [16].) The search was conducted only on graphs of order 24 to 30 to determine if a minimum order 3-e.c. graph lies in one of these two classes. We note that the vertex-transitive graphs of orders 20 to 28 were checked for the 3-e.c.







the corresponding probability space *the random Cayley graphs on the group  $H$  of order  $m$  with probability  $p$*  and write  $\mathcal{H}_m(p)$ . Random Cayley graphs have been studied especially in their connection with expansion properties of graphs; see [1].

While  $|S|$  is a random variable in  $\mathcal{H}_m(p)$ , all choices of  $S$  give rise to Cayley graphs, and hence, vertex-transitive graphs. This follows directly from the definition of Cayley graphs. We prove the following result in the case when  $p = 1/2$ . The method of proof is based on a construction of vertex-transitive  $n$ -e.c. tournaments exploiting circulant tournaments in [9].

**Theorem 4.** *For  $n$  a positive integer, a.a.s.  $\mathcal{H}_m(1/2)$  is  $n$ -e.c.*

**Proof.** Consider the graph  $G = \mathcal{H}_m(1/2)$  with order  $m$ . Fix

$$X = \{x_1, x_2, \dots, x_n\}$$

an  $n$ -set of vertices of  $G$ . We need to find a vertex  $z$  correctly joined to  $X$  (regardless of the partition of  $X$  into two sets, say  $A$  and  $B$ ). For  $v \in G$  define

$$\sigma_X(v) = \{u \in S : \text{for some } x \in X, u = vx^{-1} \text{ or } u = xv^{-1}\}$$

We would like to show we can construct a set  $U$ , disjoint from  $X$ , such that with probability tending to 1, there is a  $z \in U$  that is correctly joined to  $X$ . Equivalently, we show that with probability  $o(1)$ , there is no vertex in  $U$  correctly joined to  $X$ . We construct  $U$  such that  $|U| = \lfloor \frac{m}{4n^2} \rfloor$ , and impose the following restrictions on  $U$ .

1. For all distinct  $z$  and  $z'$  in  $U$ ,  $\sigma_X(z) \cap \sigma_X(z') = \emptyset$ .
2.  $|\sigma_X(z)| = n$ .

From the definition of  $\sigma_X$ , it is straightforward to see that item (1) ensures the event that  $z$  is joined to a vertex  $x_i$  in  $X$  is independent of the event that  $z'$  is joined to  $x_i$ , and item (2) ensures that the events that  $z$  is joined to any particular  $x_i$  are mutually independent.

We inductively construct the set  $U_k$  whose union will be  $U$ . We choose  $U_1$  to be a single vertex  $z_1$  not in  $X$  with the property that  $|\sigma_X(z_1)| = n$ . We therefore eliminate elements in  $X$  and those  $z_1$  such that  $|\sigma_X(z_1)| < n$ . For example, if it happens that  $z_1 x_i^{-1} = x_j z_1^{-1}$  for some  $i$  and  $j$ , then we must eliminate  $z_1$  from consideration. Each distinct pair of vertices from  $X$  eliminates at most one element of  $G$ . We may now find a suitable  $z_1$  since

$$m - n - \binom{n}{2} > 0.$$

(Recall that  $n$  is a constant that does not depend on  $m$ .)

Suppose that  $U_k$  has been constructed for a fixed  $k < \lfloor \frac{m}{4n^2} \rfloor$ , so that  $|U_k| = k$ , and the set  $U_k$  has elements satisfying items (1) and (2). Set  $U_k = \{z_1, \dots, z_k\}$ . We choose  $z_{k+1}$  as the new element of  $U_k$  by eliminating elements from  $V(G) \setminus U_k$ . As in the base step, by considering all the pairs of vertices from  $X$ ,  $\binom{n}{2}$  vertices are eliminated. Each vertex  $z \in U_k$  satisfies  $|\sigma_X z| = n$ . To ensure that  $\sigma_X(z) \cap \sigma_X(z') = \emptyset$  for  $z \in U_k$  and  $z' \in U_{k+1}$ , we must eliminate another  $2kn$  vertices. For large  $m$ , we may find a suitable  $z_{k+1}$  since

$$\begin{aligned} m - n - k - \binom{n}{2} - 2kn &> m \left(1 - \frac{1}{4n^2} - \frac{1}{2n}\right) - n - n^2 \\ &> 0. \end{aligned}$$

Add  $z_{k+1}$  to  $U_k$ , to form  $U_{k+1}$ . Define  $U = U_{\lfloor \frac{m}{4n^2} \rfloor}$ , so  $|U| = \lfloor \frac{m}{4n^2} \rfloor$  as desired.

We now estimate the probability that none of the vertices of  $U$  are correctly joined to  $X$ , and show this tends to 0 as  $m$  tends to  $\infty$ . By items (1) and (2), we have that

$$\mathbb{P}(\text{no } z \text{ in } U \text{ is c.j. to } X) = \left(1 - \frac{1}{2^n}\right)^{\lfloor \frac{m}{4n^2} \rfloor}.$$

Hence, we have that the probability  $P$  of the event that  $G$  is not  $n$ -e.c. satisfies

$$\begin{aligned} P &\leq \binom{m}{n} 2^n \left(1 - \frac{1}{2^n}\right)^{\frac{m}{4n^2}} \\ &\leq m^n 2^n \left(1 - \frac{1}{2^n}\right)^{\frac{m}{4n^2}} \\ &= \exp\left(n \log m + n \log 2 + \left(\frac{m}{4n^2}\right) \log\left(1 - \frac{1}{2^n}\right)\right) \\ &= o(1). \quad \square \end{aligned}$$

We note that the proof of Theorem 4 generalizes to  $p \in (0, 1)$ , and we may allow  $n$  to grow as a function of  $m$ . We omit these generalizations, as our more modest goal here is to provide a new randomized construction of vertex-transitive  $n$ -e.c. graphs.

The proof of Theorem 4 gives an asymptotic upper bound for  $\mathcal{G}_m(1/2)$  to be  $n$ -e.c., whose proof is similar to Theorem 2 and so is omitted.

**Theorem 5.** *If  $m = O(n^3 2^n)$  and  $n$  is a sufficiently large integer, then with positive probability  $\mathcal{G}_m(1/2)$  is  $n$ -e.c. In particular, there is a vertex-transitive  $n$ -e.c. graph of order  $O(n^3 2^n)$ .*

**4. Future work.** We know little about the minimum order  $n$ -e.c. graphs; for example, are they all vertex-transitive or even regular? The determination of  $m_{ec}(3)$  will likely use a mixture of computational and theoretical results on 3-e.c. graphs. By the results in [15] and those referenced in Section 2, the only orders where we do not know whether a 3-e.c. graph exists are:

$$24, 25, 26, 27, 31, 33.$$

Determining the exact order of  $m_{ec}(n)$  for  $n \geq 4$  appears to be a difficult problem. Even determining the asymptotic order of this function presents a challenge. The conjecture of Erdős that

$$m_{ec}(n) = O(n2^n),$$

remains as one of the deepest problems in this area of graph theory. Even a seemingly modest improvement to  $O(n^{2-c}2^n)$ , where  $c > 0$ , would represent a breakthrough. It is possible that random graphs stemming from either combinatorial designs (see [3]) or Cayley graphs may eventually be adapted to solve the conjecture.

## REFERENCES

1. A. Alon and Y. Roichman, *Random cayley graphs and expanders*, Random Structures and Algorithms **5** (1994), 271–284.
2. W. Ananchuen and L. Caccetta, *On the adjacency properties of Paley graphs*, Networks **23** (1993), 227–227.
3. C.A. Baker, A. Bonato, N.A. McKay, and P. Prałat, *Graphs with the  $n$ -e.c. adjacency property constructed from resolvable designs*, Journal of Combinatorial Designs **17** (2009), 294–306.
4. A. Blass, G. Exoo, and F. Harary, *Paley graphs satisfy all first-order adjacency axioms*, Journal of Graph Theory **5** (1981), 435–439.
5. B. Bollobás, *Random Graphs*, Cambridge University Press, Cambridge (2001).
6. B. Bollobás and A. Thomason, *Graphs which contain all small graphs*, European Journal of Combinatorics **2** (1981), 13–15.
7. A. Bonato, *The search for  $n$ -e.c. graphs*, Contributions to Discrete Mathematics **4** (2009), 40–53.
8. A. Bonato and K. Cameron, *On an adjacency property of almost all graphs*, Discrete Mathematics **231** (2001), 103–119.
9. A. Bonato, P. Gordinowicz, and P. Prałat, *Bounds and constructions for  $n$ -e.c. tournaments*, Preprint, 2010.
10. L. Caccetta, P. Erdős, and K. Vijayan, *A property of random graphs*, Ars Combinatoria **19** (1985), 287–294.
11. A Costea, *Computational and theoretical aspects of  $n$ -e.c. graphs*, Master’s Thesis, Wilfrid Laurier University, 2010.
12. P. Erdős and A. Rényi, *Asymmetric graphs*, Acta Mathematica Academiae Scientiarum Hungaricae **14** (1963), 295–315.
13. C.D. Godsil and G. Royle, *Algebraic Graph Theory*, Springer, New York, 2001.

14. P. Gordinowicz and P. Prałat, *The search for the smallest 3-e.c. graphs*, Journal of Combinatorial Mathematics and Combinatorial Computing **74** (2010), 129–142.
15. O. Pikhurko and M. Singh, *Extremal existentially closed graphs*, Preprint, 2011.
16. G. Royle, *Transitive graphs*, <http://units.maths.uwa.edu.au/~gordon/remote/trans/index.html>.
17. T. Spence, *Strongly regular graphs on at most 64 vertices*, <http://www.maths.gla.ac.uk/~es/srgraphs.html>.

Received February 8, 2010.

DEPARTMENT OF MATHEMATICS, RYERSON UNIVERSITY, TORONTO, ON, CANADA,  
M5B 2K3

*E-mail address:* abonato@ryerson.ca

DEPARTMENT OF MATHEMATICS, WILFRID LAURIER UNIVERSITY, WATERLOO,  
NS, CANADA, N2L 3C5

*E-mail address:* ale.costea@gmail.com